



## Gate-dependent spin-orbit coupling in multielectron carbon nanotubes

Jespersen, Thomas Sand; Grove-Rasmussen, Kasper; Paaske, Jens; Muraki, K.; Fujisawa, T.; Nygård, Jesper; Flensberg, Karsten

*Published in:*  
Nature Physics

*DOI:*  
[10.1038/NPHYS1880](https://doi.org/10.1038/NPHYS1880)

*Publication date:*  
2011

*Document version*  
Early version, also known as pre-print

*Citation for published version (APA):*  
Jespersen, T. S., Grove-Rasmussen, K., Paaske, J., Muraki, K., Fujisawa, T., Nygård, J., & Flensberg, K. (2011). Gate-dependent spin-orbit coupling in multielectron carbon nanotubes. *Nature Physics*, 7(4), 348-353. <https://doi.org/10.1038/NPHYS1880>

# Gate-dependent orbital magnetic moment in carbon nanotubes

## Supporting online material

T. S. Jespersen<sup>1\*†</sup>, K. Grove-Rasmussen<sup>1,2\*†</sup>, K. Flensberg<sup>1</sup>, J. Paaske<sup>1</sup>,  
K. Muraki<sup>2</sup>, T. Fujisawa<sup>3</sup>, and J. Nygård<sup>1</sup>.

<sup>1</sup>Niels Bohr Institute & Nano-Science Center, University of Copenhagen,  
Universitetsparken 5, DK-2100 Copenhagen, Denmark

<sup>2</sup> NTT Basic Research Laboratories, NTT Corporation,  
3-1 Morinosato-Wakamiya, Atsugi 243-0198, Japan

<sup>3</sup> Research Center for Low Temperature Physics, Tokyo Institute of Technology,  
Ookayama, Meguro, Tokyo 152-8551, Japan

<sup>†</sup> Equal contribution

## Estimating the band gap

The nanotube sample studied in the article is particularly regular in the many-electron regime (see data in article), while less so close to the band gap. The strong/weak coupling for shells in the valence/conduction band [1] indicates that the nanotube is a small band gap semiconducting nanotube [2]. Figure S1 shows the stability diagram in the region of the largest Coulomb peak spacings and the lowest currents. The very large diamond at  $V_g \sim 0.8$  V is interpreted as a result of the band-gap energy adding to the addition energy  $\sim 50 - 60$  meV. To estimate the band-gap we subtract the average charging energy and level spacings for the first hole and first electron[3] *i.e.*, the average height of the neighboring diamonds ( $\sim 25$  meV) yielding  $E_g \sim 30$  meV. The band gap parameter appearing in the article[1] is defined as  $\Delta_g = E_g/2 \sim 15$  meV, *i.e.* consistent with the value from the fit of  $g_{\text{orb}}(V_g)$  to Eq. 3 (Fig. 4 of the main manuscript).

## Additional data

To supplement the excited state spectroscopy measurements of  $g_{\text{orb}}$  discussed in the article, the device was cooled again to 4.2 K and following the method of Ref. [4],  $g_{\text{orb}}$  values were extracted from the evolution of zero-bias Coulomb blockade peaks in a parallel magnetic field. Representative data is shown in Fig.




Figure S1: Stability diagram in the  $V_g$ -region of the band gap. A low current signal is observed at the position of the arrow leading to the diamond structure indicated by the lines.

S2 for two different  $V_g$ -regions; close to the gap (top) and for many electrons (bottom). The gate axis has been converted into energy by the  $\alpha$ -factor which is measured from the corresponding stability diagrams as the ratio of Coulomb diamond heights (bias) to widths (gate). The orbital  $g$ -factor is extracted from the  $B_{||}$ -dependence of the peak positions in Fig. S2 and displayed in Fig. 4 of the main manuscript (square data points). As discussed above, the regular 4-fold periodicity is absent in the vicinity of the gap impeding direct comparison to the 4-fold model. Instead  $g_{\text{orb}}$  was measured for all charge-states in the gate region indicated by the horizontal error bars in Fig. 4 of the main manuscript and the spread in resulting  $g_{\text{orb}}$ -values is indicated by the vertical error bars.

## References

- [1] T. S. Jespersen, *et al.*, Nature Phys. **7**, 348 (2011).
- [2] J. Cao, Q. Wang, and H. Dai, Nat. Mat. **4**, 745 (2005).
- [3] F. Kuemmeth, *et al.*, Nature **452**, 448 (2008).
- [4] E. Minot, *et al.*, Nature **428**, 536 (2004).




Figure S2: The evolution of Coulomb peaks in a parallel magnetic field for two different gate regions. Close to the gap (top) and in the many electron regime (bottom). The gate axis has been converted to energy by the  $\alpha$ -factor which is extracted from the corresponding stability diagrams (see text). The  $B_{\parallel}$ -dependence of the peak positions yields  $g_{\text{orb}}$ .